

# The nature of most probable paths at finite temperatures

Pratip Bhattacharyya\*

*Theoretical Condensed Matter Physics Division, Saha Institute of Nuclear Physics,  
Sector - 1, Block - AF, Bidhannagar, Kolkata 700 064, India*

(Dated: March 28, 2003)

We determine the most probable length of paths at finite temperatures, with a preassigned end-to-end distance and a unit of energy assigned to every step on a  $D$ -dimensional hypercubic lattice. The asymptotic form of the most probable path-length shows a transition from the directed walk nature at low temperatures to the random walk nature as the temperature is raised to a critical value  $T_c$ . We find  $T_c = 1/(\ln 2 + \ln D)$ . Below  $T_c$  the most probable path-length shows a crossover from the random walk nature for small end-to-end distance to the directed walk nature for large end-to-end distance; the crossover length diverges as the temperature approaches  $T_c$ . For every temperature above  $T_c$  we find that there is a maximum end-to-end distance beyond which a most probable path-length does not exist.

PACS numbers: 05.40.Fb, 05.70.Jk

Among the different paths that can be drawn on a lattice, two of the most common are generated by random walks and directed walks. The length  $n$  of a path is measured as the total number of steps from one end to the other, where each step covers a unit lattice bond. The paths are classified according to the relation between their lengths  $n$  and the distance  $r$  between their end-points. Paths generated by a random walk are characterised by  $n \propto r^2$  [1, 2, 3] whereas paths generated by a directed walk have  $n \propto r$  [4]. The problem of determining the relation between  $n$  and  $r$  under a given condition (such as the minimum energy condition for the path) can be defined in two ways : either to determine the end-to-end distance  $r$  for paths of a preassigned length  $n$ , or, to determine the length  $n$  of paths with a preassigned end-to-end distance  $r$ . The problem of the first type is realised in linear polymers seeking a minimum energy configuration on a substrate on in a solvent [5, 6, 7], whereas the problem of the second type is only of geometric interest.

In this paper we address the problem of the second type. Consider a  $D$ -dimensional hypercubic lattice. Let us assume that every step on the lattice (i.e., to travel along one lattice bond) requires a unit of energy. Now consider two points on the lattice separated by a distance  $r$  :

$$r^2 = \sum_{i=1}^D r_i^2, \quad (1)$$

where  $r_i$  is the component of  $\mathbf{r}$  in the  $i$ -th dimension. The length  $n$  of any path connecting these two points may be written as :

$$n = \sum_{i=1}^D n_i, \quad (2)$$

where  $n_i$  is the number of lattice bonds along the path in the  $i$ -th dimension. The paths of minimum energy connecting the two points are the paths of shortest length between the two points :

$$n_{\min} = \sum_{i=1}^D |r_i|. \quad (3)$$

This may be considered as the most probable path-length at zero temperature. Now our aim is to determine the most probable path-length  $n_{\text{mp}}(r, T)$  at finite temperatures between the two given points separated by a distance  $r$ . Temperature represents noise. At  $T = 0$  there is no noise; therefore the only possible path-length between the two given points is that of the shortest course from one point to the other. At finite temperatures noise raises the value the most probable path-length above the minimum. The most probable paths at finite temperatures are not the minimum energy paths, but are paths of minimum free energy.

We first calculate the number  $N(n, r)$  of distinct paths of length  $n$  on the lattice, connecting two points separated by a distance  $r$ . The number of distinct ways of assigning  $n_i$  steps to the  $i$ -th dimension when  $n_1, \dots, n_{i-1}$  steps have been already assigned to  $i-1$  other dimensions is given by the binomial coefficient

$$\binom{n + n_i - \sum_{k=1}^i n_k}{n_i}. \quad (4)$$

Thus the number of distinct ways of assigning  $n_1, \dots, n_D$  steps to dimensions 1,  $\dots$ ,  $D$  respectively is given by the polynomial coefficient :

$$\prod_{i=1}^D \binom{n + n_i - \sum_{k=1}^i n_k}{n_i} = \frac{n!}{\prod_{i=1}^D n_i!}, \quad (5)$$

and the number of distinct ways in which  $n_i$  steps can be distributed among the positive and the negative directions in the  $i$ -th dimension such that the difference

---

\*Electronic address: pratip@cmp.saha.ernet.in

between their numbers is  $r_i$  is given by the binomial coefficient

$$\binom{n_i}{\frac{n_i - r_i}{2}}. \quad (6)$$

Therefore

$$\begin{aligned} N(n, r) &= \sum_{\{n_i\}} \frac{n!}{\prod_{i=1}^D n_i!} \prod_{i=1}^D \binom{n_i}{\frac{n_i - r_i}{2}} \\ &= \sum_{\{n_i\}} \frac{n!}{\prod_{i=1}^D \left(\frac{n_i - r_i}{2}\right)! \left(\frac{n_i + r_i}{2}\right)!}. \end{aligned} \quad (7)$$

The final expression in the above equation is obtained by expanding the binomial coefficients. The summation indices  $\{n_i\}$  are subject to the conditions of Eq. (2) and Eq. (3).

At a finite temperature  $T$  the statistical weight of a step along any path is  $\exp(-1/T)$ , since a step on the lattice requires a unit of energy. Therefore the partition function for paths of length  $n$  and end-to-end distance  $r$  is given by :

$$G_T(n, r) = N(n, r) \exp(-n/T), \quad (8)$$

and the partition function for all paths with a preassigned end-to-end distance  $r$  at the temperature  $T$  is given by :

$$G_T(r) = \sum_{n=n_{\min}}^{\infty} G_T(n, r). \quad (9)$$

For large path-lengths  $n$  and  $n_i \gg r_i$ , we can use Stirling's approximation [8] for the factorials in Eq. (7) :

$$n! \sim \sqrt{2\pi n} n^n \exp(-n), \quad (10)$$

$$\begin{aligned} \left(\frac{n_i \pm r_i}{2}\right)! &\sim \sqrt{2\pi \left(\frac{n_i \pm r_i}{2}\right)} \left(\frac{n_i \pm r_i}{2}\right)^{\left(\frac{n_i \pm r_i}{2}\right)} \\ &\times \exp\left(-\frac{n_i \pm r_i}{2}\right). \end{aligned} \quad (11)$$

Consequently Eq. (8) may be written as :

$$\begin{aligned} G_T(n, r) &\sim \frac{1}{(2\pi)^{D-1/2}} \sum_{\{n_i\}} \exp \left[ n \ln n - \sum_{i=1}^D n_i \ln n_i \right. \\ &\quad \left. + \frac{1}{2} \ln n - \sum_{i=1}^D \ln n_i + (n + D) \ln 2 \right. \\ &\quad \left. - \sum_{i=1}^D \frac{r_i^2}{2n_i} - \frac{n}{T} \right]. \end{aligned} \quad (12)$$

For large path-lengths  $n$ , the most probable value of  $n_i$ , as well as the mean of  $n_i$ , is equal to  $n/D$  for all dimensions  $i$ . Hence we assume :

$$n_i \approx \langle n_i \rangle = \frac{n}{D}. \quad (13)$$

The partition function in Eq. (12) now reduces to the form :

$$\begin{aligned} G_T(n, r) &\sim \frac{1}{(2\pi)^{D-1/2}} \exp \left[ - \left( D - \frac{1}{2} \right) \ln n \right. \\ &\quad \left. + (n + D)(\ln 2 + \ln D) \right. \\ &\quad \left. - \frac{D}{2n} r^2 - \frac{n}{T} \right]. \end{aligned} \quad (14)$$

For the most probable value  $n_{\text{mp}}$  of the path-length at temperature  $T$ , the free energy  $-T \ln G_T(n, r)$  is minimum :

$$\left[ -T \frac{d}{dn} \ln G_T(n, r) \right]_{n=n_{\text{mp}}} = 0, \quad (15)$$

which leads to the following condition :

$$2 \left[ (\ln 2 + \ln D) - \frac{1}{T} \right] n_{\text{mp}}^2 - (2D-1)n_{\text{mp}} + Dr^2 = 0. \quad (16)$$

Solving the above equation we obtain the most probable path-length as a function of the end-to-end distance and the temperature :

$$n_{\text{mp}}(r, T) = \frac{1 - \sqrt{1 - \frac{8D}{(2D-1)^2} \left( \frac{1}{T_c} - \frac{1}{T} \right) r^2}}{\frac{4}{2D-1} \left( \frac{1}{T_c} - \frac{1}{T} \right)} \quad (17)$$

where

$$\frac{1}{T_c} = \ln 2 + \ln D. \quad (18)$$

In Eq. (17) only the negative sign of the square-root is admissible, in order to ensure that  $n_{\text{mp}}$  is always positive. In the simplest case of a one-dimensional lattice it was found in an earlier study that  $T_c = 1/\ln 2$  [9]. The result for  $D = 1$  can now be obtained directly from the general expression of  $T_c$  given in the equation above.

At low temperatures, i.e.  $T < T_c$ , the expression for the most probable path-length in Eq. (17) may be written as :

$$n_{\text{mp}}(r, T) = \frac{2D}{2D-1} \frac{r_{\times}^2}{r^2} \left[ \sqrt{1 + \left( \frac{r}{r_{\times}} \right)^2} - 1 \right] \quad (19)$$

where

$$r_{\times}(T) = \frac{2D-1}{2\sqrt{2D}} \left( \frac{1}{T} - \frac{1}{T_c} \right)^{-1/2} \quad (20)$$

is a characteristic end-to-end distance depending only on the temperature and the dimension of the lattice. The significance of this distance will be evident from the following analysis. For end-to-end distances which are much smaller than  $r_{\times}$  the most probable path-length given in Eq. (19) reduces to the form :

$$n_{\text{mp}} \approx \frac{D}{2D-1} r^2, \quad r \ll r_{\times} \quad (21)$$

which has the feature of a random walk. For end-to-end distances which are much larger than  $r_{\times}$ , Eq. (19) reduces to the form :

$$n_{\text{mp}} \approx \sqrt{\frac{D}{2}} \left( \frac{1}{T} - \frac{1}{T_c} \right)^{-1/2} r, \quad r \gg r_{\times} \quad (22)$$

which has the feature of a directed walk. Thus  $r_{\times}(T)$  is a crossover distance which marks the crossover in the nature of the most probable path-length from the random walk class to the directed walk class at any temperature  $T < T_c$ . Eq. (20) shows that the crossover distance diverges as the temperature is raised to  $T_c$  which may be formally identified as a critical point. The asymptotic form (i.e., for large  $r$ ) of the most probable path-length undergoes a transition from the directed walk nature at  $T < T_c$  to the random walk nature at  $T_c$ . The limiting form of the most probable path-length as the temperature approaches  $T_c$  may be obtained directly from Eq. (17) :

$$n_{\text{mp}} = \frac{D}{2D-1} r^2, \quad T \rightarrow T_c. \quad (23)$$

This result for  $T \rightarrow T_c$  is true for all values of  $r$ .

For high temperatures, i.e.  $T > T_c$ , the most probable path-length in Eq. (17) may be written as :

$$n_{\text{mp}}(r, T) = \frac{2D r_{\text{max}}^2}{2D-1} \left[ 1 - \sqrt{1 - \left( \frac{r}{r_{\text{max}}} \right)^2} \right]. \quad (24)$$

where

$$r_{\text{max}}(T) = \frac{2D-1}{2\sqrt{2D}} \left( \frac{1}{T_c} - \frac{1}{T} \right)^{-1/2} \quad (25)$$

is another characteristic end-to-end distance depending only on the temperature and the dimension of the lattice. At any temperature  $T > T_c$ , Eq. (24) shows that for end-to-end distances  $r > r_{\text{max}}(T)$  a real and finite value of the most probable path-length does not exist. For end-to-end

distances much smaller than  $r_{\text{max}}(T)$  Eq. (24) reduces to the form :

$$n_{\text{mp}} \approx \frac{D}{2D-1} r^2, \quad r \ll r_{\text{max}} \quad (26)$$

which has the same random walk feature found in Eq. (21) and Eq. (23). Eq. (20) and Eq. (25) show that the two characteristic end-to-end distances below and above  $T_c$ , i.e.  $r_{\times}$  and  $r_{\text{max}}$ , diverge in the same way as  $T \rightarrow T_c$ .

In this paper we have studied a temperature driven transition in the geometry of most probable paths on a  $D$ -dimensional hypercubic lattice. In the low temperature phase ( $T < T_c$ ) the most probable paths show the appearance of a random walk for small end-to-end distances but eventually cross over to the form of a directed walk for large end-to-end distances. Close to the critical temperature ( $T \rightarrow T_c$ ) the most probable paths are like random walks for all end-to-end distances. In the high temperature phase ( $T > T_c$ ) the very existence of most probable paths is restricted by a maximum end-to-end distance for each temperature. The critical temperature is found to be  $T_c = 1/(\ln 2 + \ln D)$  which gets lower with increasing dimension of the lattice because of a simple reason : the larger the dimension, the greater the entropy of the paths, and therefore, the lower is the noise (given as the temperature) required to bring the random walk nature.

### Acknowledgments

I am grateful to Bikas K. Chakrabarti and D. Stauffer for their comments on the manuscript.

- 
- [1] S. Chandrasekhar, Rev. Mod. Phys. **15**, 1 (1943).
  - [2] F. Reif, *Fundamentals of Statistical and Thermal Physics*, (McGraw-Hill, New York, 1965).
  - [3] J. W. Haus and K. W. Kehr, Phys. Rep. **150**, 263 (1987).
  - [4] T. Halpin-Healy and Y-C. Zhang, Phys. Rep. **254**, 215 (1995).
  - [5] P-G. deGennes, *Scaling Concepts in Polymer Physics*, (Cornell University Press, Ithaca, 1979).
  - [6] K. Barat and B. K. Chakrabarti, Phys Rep. **258**, 377 (1995).
  - [7] T. A. Witten, Rev. Mod. Phys. **70**, 1531 (1998).
  - [8] R. L. Graham, D. E. Knuth and O. Patashnik, *Concrete Mathematics*, (Addison-Wesley, 2nd ed., 1994).
  - [9] P. Bhattacharyya, Y. M. Strelniker, S. Havlin and D. ben-Avraham, Physica A **297**, 401 (2001).